

**Final Report
on
NASA Grant NAGW-1101**

**Extended Ecosystem Signatures with Application
to Eos Synergism Requirements**

(NASA-CR-193398) EXTENDED
ECOSYSTEM SIGNATURES WITH
APPLICATION TO EOS SYNERGISM
REQUIREMENTS Final Report
(Michigan Univ.) 24 p

N94-11374

Unclas

G3/45 0176677

July 1993



THE UNIVERSITY OF MICHIGAN

**Radiation Laboratory
Department of Electrical Engineering
and Computer Science
Ann Arbor, Michigan 48109-2122
USA**

Technical Report 024557-1-F

**Final Report
on
NASA Grant NAGW-1101**

**Extended Ecosystem Signatures with Application to
Eos Synergism Requirements**

July 1993

**Principal Investigator: Fawwaz T. Ulaby, University of Michigan
Co-Investigators: M. Craig Dobson, University of Michigan
Kamal Sarabandi, University of Michigan**

Table of Contents

1.0 Introduction

2.0 Synopsis of Research Results

2.1 Ground-Based Experiments

2.2 Model Development and Validation

2.3 Airborne Experiments

2.4 Extension of MIMICS to Open Canopies

2.5 MIMICS Augmentations

2.6 Model Inversions

2.7 User Friendly MIMICS Code

3.0 List of Publications and Presentations

1.0 Introduction

The primary objective of this study is to define the advantages of synergistically combining optical and microwave remote sensing measurements for the determination of biophysical properties important in ecosystem modeling. This objective has been approached in a stepwise fashion starting with ground-based observations of controlled agricultural and orchard canopies and progressing to airborne observations of more natural forest ecosystems. This observational program is complemented by a parallel effort to model the visible reflectance and microwave scattering properties of composite vegetation canopies. The goals of the modeling studies are to verify our basic understanding of the sensor-scene interaction physics and to provide the basis for development of inverse models optimized for retrieval of key biophysical properties. These retrieval algorithms can then be used to simulate the expected performance of various aspects of Eos including (1) the need for simultaneous SAR and HIRIS observations or justification for other (non-synchronous) relative timing constraints and (2) the frequency, polarization and angle of incidence requirements for accurate biophysical parameter extractions.

This final report covers the period of performance from June 1987 to June 1993. During this period, the following institutions received support or collaborated with the University of Michigan on this effort:

- Aster Consulting
- Duke University
- Environmental Research Institute of Michigan
- Fresno State
- George Washington University
- Jet Propulsion Laboratory
- NASA Ames Research Center
- University of California at Davis
- University of Nebraska
- University of Wisconsin

Support from this project has contributed to 6 PhD degrees and funded the efforts of over 40 other graduate and undergraduate research assistants. The results of this research have been distributed via 119 papers including: theses, technical reports, symposia, and journal articles. These are listed in Section 3. In

addition, the MIMICS computer code for microwave scattering from vegetation has been presented at several workshops and installed, at no cost, at over 20 sites in the North America, Europe, Asia and Australia.

2.0 Synopsis of Research Results

This program completed a very successful series of truck-mounted experiments, made remarkable progress in development and validation of optical reflectance and microwave scattering models for vegetation, extended these results through airborne experimentation, expanded the scattering models to accommodate discontinuous and periodic canopies, developed inversion approaches for surface and canopy properties and disseminated these results widely through symposia and journal publications. In addition, the third generation of the computer code for the microwave scattering models has been provided to a number of other US, Canadian, Australian and European investigators who are currently presenting and publishing results using the MIMICS research code.

2.1 Ground-Based Experiments

A highly successful field experiment was conducted in the summer of 1987 in a walnut orchard at the Kearney Agricultural Field Station near Fresno, California. This experiment and follow-up studies have involved over 50 researchers from 11 institutions (The University of Michigan, the University of Nebraska, the University of California at Davis, the University of Wisconsin, Duke University, George Washington University, Fresno State, ERIM, the Jet Propulsion Laboratory, Ames Research Center and Aster Consulting). In this experiment, the orchard was monitored over a several week period in mid-summer by boom-mounted optical radiometers (Barnes MMR) and polarimetric microwave scatterometers (L-, C- and X-band range-probing, scatterometers). The observations were made in a continuous fashion over adjacent diurnal periods such that temporal changes in optical reflectance and/or radar backscattering could be correlated with temporal variations in canopy conditions. Key to the experiment was a thorough suite of observations of the canopy biophysical variates including: biomass (total and by canopy component such as trunks and branches), leaf area index, size and

orientation distributions of foliage and woody components, the moisture status of vegetation components (gravimetric moisture and water potential), soil properties (surface roughness, bulk density and moisture distribution), optical reflectance and transmittance properties of the soil and vegetation components, and the microwave dielectric properties of the soil and vegetation components. This experiment has now been extensively reported in the literature in terms of the objectives, data and findings.

The walnut experiment has produced a number of significant findings. Optical reflectance is shown to vary over the day, but only in response to changes in solar zenith angle. Consequently, preferred observation times for HIRIS in the optical region are not driven, in this case, by diurnal changes in canopy properties. Thus, no case can be made for simultaneous observation of vegetated terrain by HIRIS and SAR. As a result, HIRIS timing should be linked to expected minima in local cloud-cover.

In contrast, the microwave backscattering from the orchard exhibited pronounced diurnal fluctuations in response to variations of soil and canopy water status. Tests of a novel method for nondestructive, *in situ* measurement of the microwave dielectric constant of woody components of the walnut trees demonstrated that circadian patterns of water flow and distribution are the primary causative factors leading to diurnal variation in radar backscattering of 2 dB to 3 dB. An important technical result of this study is that while the signal from the Eos SAR can be expected to be diurnally variant in response to changes in canopy and substrate moisture conditions and changes in leaf angle distributions (at high frequencies) for heliotropic plants, these variations are shown to be relatively continuous in nature with certain notable exceptions. Truck-mounted scatterometer data and polarimetric AIRSAR data (from other experiments) demonstrate that certain meteorologic events (i.e., rain) and anthropogenic activities (i.e., tillage and harvest) induce an immediate change in radar backscattering.

2.2 Model Development and Validation

In addition, great progress has been made in the development and validation of models to predict radar backscatter and optical reflectance on the basis of measurable biophysical attributes. The one-dimensional Cupid model predicts bidirectional reflectance functions which agree very closely with the MMR measurements of

the walnut orchard. In the microwave region, MIMICS I, a first-order, vector radiative transfer model, is found to predict the observed L-band backscattering coefficient to within better than 1 dB and closely matches the observed diurnal variations at all polarizations. Despite this excellent agreement at L-band, some of the model assumptions begin to break down at higher frequencies (X-band); and it is apparent that single-scattering theory is insufficient to explain X-band cross-polarized backscattering in particular.

2.3 Airborne Experiments

In an effort to extend the preceding findings from an "engineered" orchard to conditions more representative of a natural forest ecosystem, a multirate airborne SAR experiment was undertaken in July through September of 1989 using the JPL AIRSAR. Approximately 20 researchers from the University of Michigan, ERIM, Duke University, JPL and the University of California at Davis collected extensive ground-level observations from 60 forest stands at the Duke University Research Forest. These even-aged stands of primarily loblolly pine are distributed over two soil types and ranged in age from seedlings to about 80 years. Microwave dielectric measurements made *in situ* in living loblolly trunks were used to investigate the variation in the dielectric constant as a function of depth into the trunk tissue, as a function of height and as a function of time over several diurnal periods. These measurements, made at P-, L-, C- and X-bands, show diurnal variations which are highly correlated with tree physiologic properties such as stem water-potential and transpiration which, in turn, are linked to cloud-cover and incident solar radiation.

In early September, the JPL AIRSAR overflew the Duke Research Forest with 12 identical passes 20 minutes apart from noon until about 4:00 pm and again the next morning with 3 passes from 8:00 am to 9:00 am. Radar backscatter simulations with MIMICS I using the measured tree structural and dielectric properties over the experimental period show a diurnal variation of generally less than 1 dB at P-, L- and C-bands for the forest stands of loblolly pine. These model expectations are congruent with externally calibrated AIRSAR observations. Since the uncertainties in mean radar backscattering coefficient related to coherent fading and calibration accuracy are both on the order of ± 1 dB, the observed temporal variation in

radar backscattering from these forest stands is not very significant. While the humid loblolly pine forest of North Carolina did not exhibit significant diurnal variation in radar backscattering coefficient, this forest was clearly not moisture-stressed. Hence, care must be taken in extrapolating these results to other forest environments. However, it is clear from the dielectric and radar evidence collected to date that, for studies seeking to maximize sensitivity to circadian patterns in plant water status, Eos SAR passes close to (or shortly after) solar noon and again pre-dawn would be the preferred times.

Recent comparisons of calibrated backscattering coefficients extracted from AIRSAR data for the Duke forest loblolly pines with similar data for maritime pines of the Landes Forest near Bordeaux, France show very strong correlations with aboveground biomass at P- and L-bands, especially for H-polarized terms. The power-law relationship between backscattering and biomass is found to be identical for both species for all polarizations except at C-band (where one specie is offset by 1dB to 2 dB relative to the other). The long wavelength response is the same for both species because the trunk and branch architectures of the two species are similar. The differences at C-band are attributable to inter-specie differences in the size, orientation and density of needles. Importantly, this power-law relationship at long wavelengths is predicted by MIMICS.

2.4 Extension of MIMICS to Open Canopies

The MIMICS I model assumes a vegetation canopy to be comprised of three statistically homogeneous random layers. The scattering elements in the crown and trunk layers are uniformly distributed over a dielectric half-space with a rough surface. As a consequence, the model is only applicable to "closed-canopy" conditions meaning there are no gaps in the crown. The crown constituents, foliage and branches, are modeled as dielectric discs or needles and smooth dielectric cylinders, respectively. The trunk layer is assumed to consist of a size distribution of homogeneous dielectric cylinders. The air-crown and crown-trunk interfaces are treated as diffuse boundaries. Since the relative volume fraction of canopy elements is low ($\ll 1\%$), the phase and extinction matrices in the radiative transfer formulation are derived using the single scattering approximation.

The "closed-canopy" assumption of MIMICS I is often violated in natural ecosystems where there may be gaps in the crown due to

mortality and uneven seedling establishment or due to a mix of plant species with very different growth habit. In order to generalize MIMICS, we have developed MIMICS II which treats the case of a discontinuous crown layer. MIMICS II considers two sets of random variables: (1) tree-level random variables which characterize the scattering elements within the crown section of individual trees and (2) canopy-level random variables which characterize the variability among trees (such as the heights of trees, the sizes and shapes of crown sections for each tree, and the locational distribution of trees). Tests of MIMICS II show that, for forest stands characterized by a crown layer with low particle density (that is few branches and leaves), the model yields results not significantly different than those of MIMICS I. As the particle density within individual crowns increases, the two models diverge especially at high frequencies. At low frequencies (such as P-band) and incidence angles away from nadir, even trees with well-developed crowns yield nearly identical model predictions using the continuous or discontinuous canopy formulation. Since MIMICS II is much more computationally intensive, it is important to know that the MIMICS I assumption of a "closed-canopy" is adequate at low frequencies.

Both MIMICS I and II assume that individual plants are randomly distributed. In many agricultural crops, orchards and tree plantations, the locations of individual plants are not randomized, rather there is a periodic organization in at least one lateral dimension. MIMICS III has been developed to treat these cases. It has been successfully tested against data obtained from corn fields by the University of Michigan's truck-mounted POLARSCAT during the summer of 1990.

The MIMICS models have shown remarkable success in predicting the radar response of vegetation canopies, particularly at L-band. Despite its success, MIMICS has a number of inherent limitations and deficiencies.

2.5 MIMICS Augmentations

Extension to Second Order

Whereas at L-band the backscattering contributions of the leaves and branches in the crown layer are secondary in importance in comparison to the contributions to net backscattering by the trunk

and ground layers, the leaves and branches become equally important, and sometimes dominant contributors, at higher frequencies (X-band). Hence, at L-band it is sufficient to account for only first-order scattering by the leaves and branches, but this is not the case at X-band, particularly for the cross-polarized component of the backscattering coefficient. This is evidenced by the poor agreement between the MIMICS I predictions of the cross-polarized response and experimental data for the walnut orchard at X-band. In part, this is a consequence of the size distributions of the foliage and branches becoming comparable to or greater than wavelength. Moreover, even at low frequencies (P- and L-band), the inclusion of second-order interactions between the trunks and the ground surface (trunk-trunk-ground, trunk-ground-trunk, and ground-trunk-trunk contributions) are found to improve the accuracy of MIMICS.

Improvement of Constituent Scattering Models

In MIMICS, as well as in all other available vegetation scattering models, the leaves are treated as flat disks and the branches as straight, smooth cylinders or ellipsoids at low frequencies. In reality, leaves are not flat, and the needles may be comparable to or longer than wavelength over the desired frequency range. Models have been developed for both curved leaves and long needles. Results show the reduction in radar cross section is a function of the radius of curvature relative to wavelength.

A model has also been developed for scattering by a multi-layered rough dielectric cylinder which is a more realistic representation of the dielectrically non-homogeneous bark and inner core of trunks and major branches. For the cases tested to date, the use of a two-layer corrugated cylinder model leads to a reduction in radar cross-section.

In many cases, dihedral-like multiple specular scattering mechanisms (such as ground-trunk) dominate the net backscatter from the forest. This is particularly true at long wavelengths. Laboratory based scatterometer measurements verify the importance of the roughness of the surface layer in controlling specular scattering. This roughness dependence is modeled and incorporated into MIMICS.

Derivation of Phase Statistics from Canopy Models

Since phase data is available directly from the output of polarimetric SARs, the information contained in the relative phase differences between polarization states can be used to infer information about the canopy. Because scattering models based upon radiative transfer do not provide phase information directly, it has not been possible to compare distributions of phase data to theoretical expectations. A technique has been developed that provides the phase difference statistics completely from the averaged Mueller matrix produced by the scattering model. Examination of the phase difference statistics obtained from the canopy models based on this technique compare favorably with scatterometer and AirSAR observations.

2.6 Model Inversions

Since canopy models, such as MIMICS, are usually complicated and highly non-linear functions of the desired biophysical parameters, direct inversion of these models are not straightforward. A standard approach for inversion of such models is an optimization technique that starts with an initial estimate of scene parameters and searches for convergence upon an acceptable solution via an iterative process. Such optimization techniques require calculation of the direct scattering problem many times before converging upon the solution. Because of the complexity of the vegetation scattering models, this is a computationally inefficient process. Three alternative approaches have been investigated: (1) artificial neural networks, (2) an empirical iterative approach and (3) a semi-empirical approach. The first two of these approaches use MIMICS to simulate a data sets over a ranges of scene conditions which are impractical to collect by airborne experimentation. These simulated data sets contain subsets which cover the available airborne SAR observations.

Artificial Neural Network Approach

In the past few years, the artificial neural network has been applied to several types of remote sensing problems. Neural networks offer three major advantaged over other inversion approaches. The first is the massive parallelism used in neural networks which provides higher computational rates than can be provided by serial computation. The second advantage is that the algorithms for hte neural network are very general and cna be used

as a "black box" for any desired model or sets of input-output data. The third advantage is that neural network software is widely available.

Initial attempts to train a neural network using actual SAR data for the Duke Forest data set met with only limited success. We suspect that this is in part due to the narrow range of scene conditions present within that data set. As a consequence, we trained the neural network using data simulated by MIMICS over a much broader range of conditions using the backpropagation learning algorithm. The qualitative and quantitative performance of the network is found to yield promising results for the determination of forest biophysical properties such as biomass, stem density, basal area and height.

Empirical Iterative Approach

A canopy model can be viewed as a mult-valued, nonlinear continuous vector function. The choice of the output vector, and therefore the function itself, is at the discretion of the system designer, while the variables of the function and their domain is specified by the canopy configuration. The quantities measured by a polarimetric SAR are the amplitude and phase of the two co-polarized and two cross-polarized channels for any desired number of frequencies and incidence angles. Frequency and incidence angle must be selected such that the scattering model shows the maximum sensitivity to the parameters of interest. If the function is one-to-one over the desired domain of the input parameters, then invertibility and uniqueness of the inverse solution are guaranteed. The canopy models are not, in general, one-to-one functions; however, they are usually very smooth functions of the input parameters and do not show any strong resonance behavior. Consequently, the parameter space may be divided into some finite subspaces for which the function is one-to-one. Another difficulty in finding the inverse function is that the function is nonlinear in most cases. A linear system of equations is the only vector function that can be inverted exactly and systematically. It would then be desirable to find some transformation of the system response into a region over which the behavior is approximately linear. This forms the basis of the iterative inversion algorithm which works with the piecewise linear function.

For polarimetric SAR data available over a range of incidence angles, the data may be represented in terms of a Fourier series in the restricted angular range. The behavior of the Fourier coefficients

as a function of the canopy parameters provide a convenient measure of the system response that is independent of incidence angle. Since the angular dependency of radar backscatter away from nadir is smooth, only the first few terms in the Fourier series are needed to represent the function. If the behavior of each Fourier component is known in terms of the canopy parameters, then we have an empirical model that describes the the scene behavior in detail. The functional form of the Fourier coefficients might be nonlinear which can be piecewise linearized over nested subdomains in the parameter space. The resulting sets of linear equations are directly invertible; and by means of a iterative process that successively converges on smaller domains, a solution set can be found.

Results obtained with this approach on a simple canopy model indicates that the algorithm is very efficient and fast, once the subdomains and linear functions are obtained. The advantage of this method over the neural network approach is that here a set of measurables can be chosen for inversion for which the maximum sensitivity to the desired parameters exist.

Semi-Empirical Model for Retrieval of Soil Properties

A third approach uses a semi-empirical model to estimate near-surface soil moisture and surface roughness from multi-polarized radar backscatter. The model is based upon backscatter data obtained from bare soil surfaces by truck-mounted polarimetric scatterometers. The approach uses polarization ratios in a semi-empirical formulation to yield estimates of soil dielectric properties and RMS roughness of the air/soil interface. Soil moisture is inferred from the estimate of the relative dielectric constant. The model performs very well for scatterometer data and is in the process of being validated for airborne SAR data acquired in Michigan, Oklahoma and California in collaboration with NASA Goddard Space Flight Center and the Jet Propulsion Laboratory.

2.7 User Friendly MIMICS Code

MIMICS has been distributed *gratis* to a large number of research institutions including: the Jet Propulsion Laboratory, the Canadian Center for Remote Sensing, the University of Guelph, Laval University, Goddard Space Flight Center, the Environmental Research

Institute of Michigan, Paul Sabatier University, Stanford University, the Joint Research Center of the EC at Ispra, and the University of Stuttgart among others. As initially distributed, MIMICS is a research code which contains a library of numeric computational modules describing the various scattering models for surfaces and vegetation. However, as a research code, these numeric modules are somewhat incompatible and not at all "user-friendly". The modules do not take advantage of recent developments in parallel computation, computer network technology, or artificial intelligence. As a result, it was felt that many potential academic researchers could not take full advantage of MIMICS (I to III).

A portable, integrated user interface and analysis package was developed for the MIMICS models. This package is now available for wider distribution to other institutions interested in incorporating the software into their educational or research activities. This package is highly portable to all UNIX-based platforms, and employs a variety of software components for analysis, inference and visualization of data.

3.0 List of Publications and Presentations

- [1] Cimino, J. B., C. Bruegge, D. Diner, J. Paris, M. C. Dobson, D. Gates, F. Ulaby, N. Goel, E. Kasischke, D. Kimes, R. Lang, J. Norman, and V. Vanderbilt, "Synergism Requirements and Concepts for SAR and HIRIS on Eos," IEEE International Geoscience and Remote Sensing Symposium (IGARSS '87), IGARSS '87 Digest, IEEE-87CH2434-9, Vol. II, pp. 955-966, Ann Arbor, Michigan, May 18-21, 1987.
- [2] Sieber, A. J., J. B. Cimino, R. Brown, J. Cihlar, M. C. Dobson, J. Ford, D. Gates, P. H. Hartl, G. Hildebrandt, R. Hoffer, M. Imhoff, E. Kasischke, J. P. Malingreau, J. Megier, T. Milne, J. Paris, J. Richards, F. Rocca, B. Rock, M. Sami, D. Sikmonett, T. LeToan, and F. T. Ulaby, "The Global Forest Ecosystem as Viewed by ERS-1, SIR-C and Eos," IEEE International Geoscience and Remote Sensing Symposium (IGARSS '87), IGARSS '87 Digest, IEEE-87CH2434-9, Vol. II, pp. 967-974, Ann Arbor, Michigan, May 18-21, 1987.
- [3] EOS Synergism Study Data Report, JPL Document D-6097, Jet Propulsion Laboratory, California Institute of Technology, 1988.
- [4] Dobson, M. C., F. T. Ulaby, and J. Paris, "Radar Backscatter from Tree Canopies," Forest Signatures Workshop, Joint Research Center, Ispra, Italy, September 4-6, 1988.
- [5] Curtiss, B. and S. L. Ustin, "The Remote Detection of Early Stages of Air Pollution Injury in Coniferous Forests Using Imaging Spectrometry," Forest Signatures Workshop, Joint Research Center, Ispra, Italy, September 4-6, 1988.

- [6] Cimino, J. B., J. Paris, M. C. Dobson, D. Gates, J. A. Weber, F. T. Ulaby, S. L. Ustin, V. Vanderbilt, J. Norman, R. Lang, and E. Kasischke, "EOS Synergism Study: Overview and Objectives," International Geoscience and Remote Sensing Symposium (IGARSS '88), Edinburgh, Scotland, September 12 - 16, 1988.
- [7] Ulaby, F. T., M. W. Whitt, and M. C. Dobson, "Radar Polarimetric Observations of a Tree Canopy," International Geoscience and Remote Sensing Symposium (IGARSS '88), IGARSS'88 Digest, IEEE-88CH2497-6, Vol. 2, pp. 1005-1008, Edinburgh, Scotland, September 12 - 16, 1988.
- [8] Caldwell, W. and V. C. Vanderbilt, "Field System to Determine Reflectance of Tree Canopies," International Geoscience and Remote Sensing Symposium (IGARSS '88), Edinburgh, Scotland, September 12 - 16, 1988.
- [9] Dobson, M. C., "Diurnal and Seasonal Variations in the Microwave Dielectric Constant of Selected Trees," International Geoscience and Remote Sensing Symposium (IGARSS '88), IGARSS'88 Digest, IEEE-88CH2497-6, Vol. 3, Edinburgh, Scotland, September 12 - 16, 1988.
- [10] Ulaby, F. T., K. McDonald, K. Sarabandi, and M. C. Dobson, "Michigan Microwave Canopy Scattering Model (MIMICS)," International Geoscience and Remote Sensing Symposium (IGARSS '88), IGARSS'88 Digest, IEEE-88CH2497-6, Vol. 2, pp. 1009-1012, Edinburgh, Scotland, September 12 - 16, 1988.
- [11] Dobson, M. C., K. McDonald, F. T. Ulaby, and J. F. Paris, "Diurnal Patterns in Multifrequency, Multipolarization Backscattering by a Walnut Orchard," International Geoscience and Remote Sensing Symposium (IGARSS '88), Edinburgh, Scotland, September 12 - 16, 1988.
- [12] Ustin, S. L., S. N. Martens, J. M. Norman, and D. Goldhammer, "Measurement and Characterization of Tree Canopy Architecture," International Geoscience and Remote Sensing Symposium (IGARSS '88), IGARSS'88 Digest, IEEE-88CH2497-6, Vol. 3, pp. 1753, Edinburgh, Scotland, September 12 - 16, 1988.
- [13] McDonald, K., M. C. Dobson, and F. T. Ulaby, "Determination of Soil Moisture Beneath a Stalk or Trunk Dominated Canopy by Radar," International Geoscience and Remote Sensing Symposium (IGARSS '88), Edinburgh, Scotland, September 12- 16, 1988.
- [14] Weber, J. A. and S. L. Ustin, "Water Relations of a Walnut Orchard: Simultaneous Measurement with Remote Sensing," International Geoscience and Remote Sensing Symposium (IGARSS '88), IGARSS'88 Digest, IEEE-88CH2497-6, Vol. 3, pp. 1749-1752, Edinburgh, Scotland, September 12 - 16, 1988.
- [15] Dobson, M. C., T. Lewis, and E. Kasischke, "Modeling of Microwave Interactions with Forest Canopies", Second Annual Strategic Relocatable Target Workshop, MIT Lincoln Labs, 16-17 May, 1989.
- [16] Senior, T.B.A., and K. Sarabandi, "Scattering by a narrow gap", Proc. of IEEE Trans. Antennas Propagat., San Jose, June 1989.

- [17] Cimino, J., J. Paris, C. Dobson, F. Ulaby, J. Weber, V. Vanderbilt, S. Ustin, J. Norman, R. Lang, E. Kasischke, and Ray Hunt, "Synergism Study: Implications of Diurnal Change in Vegetation Canopies on Eos Orbit Selection," 1989 International Geoscience and Remote Sensing Symposium (IGARSS '89), IGARSS'89 Digest, IEEE-89CH2768-0, Vol. 3, pp. 1365, July 10-14, 1989, Vancouver, Canada.
- [18] Dobson, M. C. and E. S. Kasischke, "Microwave Attenuation by Boreal Forest Canopies in Winter," 1989 International Geoscience and Remote Sensing Symposium (IGARSS '89), July 10-14, 1989, Vancouver, Canada.
- [19] McDonald, K. C., M. C. Dobson, and F. T. Ulaby, "Using MIMICS to Model Microwave Backscatter from Tree Canopies," 1989 International Geoscience and Remote Sensing Symposium (IGARSS '89), IGARSS'89 Digest, IEEE-89CH2768-0, Vol. 4, pp.2491, July 10-14, 1989, Vancouver, Canada.
- [20] Whitt, M. W., F. T. Ulaby, and M. C. Dobson, "Polarimetric Measurement of the Propagation Characteristics of Forest Clutter," 1989 International Symposium on Noise and Clutter Rejection in Radars and Imaging Sensors, November 14-16, 1989, Kyoto, Japan.
- [21] Caldwell, W. and V. C. Vanderbilt, "Canopy Radiance Measurement System," Optical Engineering, Vol. 28, pp. 1227-1236, 1989.
- [22] Curtiss, B. and S. L. Ustin, "The Remote Detection of Early Stages of Air Pollution Injury in Coniferous Forests Using Imaging Spectrometer," accepted by International Journal of Remote Sensing, 1989.
- [23] Tassoudje, A., K. Sarabandi, and F. T. Ulaby, "Design Consideration and Implementation of the LCX Polarimetric Scatterometer (POLARSCAT)," Radiation Laboratory Technical Report 022486-T-A, EECS Dept., Univ. of Michigan, Ann Arbor, Michigan, June 1989.
- [24] Sarabandi, K., F.T. Ulaby, and T.B.A. Senior, "Scattering model for a leaf at millimeter wavelengths", Radiation Laboratory Report No. 026247-1-T, The University of Michigan, June 1989.
- [25] Sarabandi, K. and F. T. Ulaby, "Scattering from Variable Resistive and Impedance Sheets," 1989 Progress in Electromagnetic Research Symposium (PIERS), July 25-26, 1989, Massachusetts Institute of Technology, Cambridge, Massachusetts.
- [26] Whitt, M. W. and F. T. Ulaby, "Bistatic Scattering from a Dielectric Cylinder Embedded in a Layer of Discrete Particles," 1989 Progress in Electromagnetic Research Symposium (PIERS), July 25-26, 1989, Massachusetts Institute of Technology, Cambridge, Massachusetts.
- [27] Sheen, D. R., A. Freeman, and E. S. Kasischke, "Phase Calibration of Polarimetric Radar Images", IEEE Transactions Geoscience and Remote Sensing, Vol. 27, No. 4, pp. 719-731, 1989.

- [28] Ustin, S. L. and S. N. Martens, "Measurement and Characterization of Tree Canopy Architecture," Bull. Ecol. Soc., 70:285, Ecological Society of America 74th Annual Meeting, Toronto, Ontario, August 6-10, 1989.
- [29] Sarabandi, K., F.T. Ulaby, and T.B.A. Senior, "Millimeter Wave Scattering Model for a Leaf", Radio Science, V. 25, N. 1, pp. 9-18, Jan.-Feb. 1990.
- [30] Sarabandi, K., "Study of periodic dielectric surfaces: Simulation with anisotropic layers and application to high frequency scattering from corrugated convex bodies.", Radiation Laboratory Report No. 024557-2-T, The University of Michigan, February 1990.
- [31] Ulaby, F. T., M. W. Whitt, and M. C. Dobson, "Measuring the Propagation Properties of a Forest Canopy Using a Polarimetric Scatterometer," IEEE Transactions on Antennas and Propagation, Vol. 38, No. 2, pp. 251-258, February, 1990.
- [32] Ulaby, F.T., T.H. Haddock, and Y. Kuga, "Measurement and Modeling of Millimeter-Wave Scattering From Tree Foliage," Radio Science, Vol. 25, pp. 193-203, 1990.
- [33] Whitt, M. and F. T. Ulaby, "Modelling Periodic Changes in Density (Row Crops) using Radiative Transfer Theory," URSI Specialist Meeting on Signature Problems in Microwave Remote Sensing, 16-18 May, 1990, Hyannis, Massachusetts.
- [34] Ulaby, F.T. and K.C. McDonald, "MIMICS Verification for a Tree Canopy at L-Band," URSI Specialist Meeting on Signature Problems in Microwave Remote Sensing, 16-18 May, 1990, Hyannis, Massachusetts.
- [35] Sarabandi, K., and T.B.A. Senior, "Low frequency scattering from cylindrical structures at oblique incidence", Proc. of International Union of Radio Science, Dallas, May 1990.
- [36] Christensen, N. L., E. S. Kasischke, and M. C. Dobson, "The Use of SAR-Derived Estimates of Aboveground Biomass in Forest Ecosystem Models", International Geoscience and Remote Sensing Symposium (IGARSS'90), IGARSS'90 Digest, IEEE-90CH2825-8, Vol. 2, pp. 1209-1212, University of Maryland, College Park, Maryland, May 20-24, 1990.
- [37] McDonald, K. and F. T. Ulaby, "MIMICS II: Radiative Transfer Modeling of Discontinuous Tree Canopies at Microwave Frequencies," International Geoscience and Remote Sensing Symposium (IGARSS'90), IEEE-90CH2825-8, University of Maryland, College Park, Maryland, May 20-24, 1990.
- [38] Ustin, S. L., V. C. Vanderbilt, J. Way, F. Ulaby, C. Dobson, and D. M. Gates, "Opportunities for EOS Instrument Synergism in Monitoring of Forest Ecosystems", International Geoscience and Remote Sensing Symposium (IGARSS'90), IGARSS'90 Digest, IEEE-90CH2825-8, Vol. 3, pp. 2301-2304, University of Maryland, College Park, Maryland, May 20-24, 1990.
- [39] Dobson, M. C., K. McDonald, F. T. Ulaby and J. Way, " Effects of Temperature on Radar Backscatter from Boreal Forests", International

Geoscience and Remote Sensing Symposium (IGARSS'90), IGARSS'90 Digest, IEEE-90CH2825-8, Vol. 3, pp. 2481-2484, University of Maryland, College Park, Maryland, May 20-24, 1990.

- [40] Kasischke, E. S., J. B. Way, N. L. Christensen, and M. C. Dobson, "Ecosystem Monitoring with the Eos SAR", International Geoscience and Remote Sensing Symposium (IGARSS'90), IGARSS'90 Digest, IEEE-90CH2825-8, Vol. 3, pp. 2297-2300, University of Maryland, College Park, Maryland, May 20-24, 1990.
- [41] Ulaby, F. T., K. Sarabandi, K. McDonald, M. Whitt, and M. C. Dobson, "Michigan Microwave Canopy Scattering Model (MIMICS)," International Journal of Remote Sensing, Vol. 11, No. 7, pp. 1223-1253, July 1990.
- [42] Kasischke, E. S. and N. L. Christensen, Jr., "Connecting Forest Ecosystem and Microwave Backscatter Models," International Journal of Remote Sensing, Vol. 11, No. 7, pp. 1277-1298, July 1990.
- [43] McDonald, K. C., M. C. Dobson, and F. T. Ulaby, "Using MIMICS to Model L-Band Multiangle and Multitemporal Backscatter for a Walnut Orchard", IEEE Transactions on Geoscience and Remote Sensing, Vol. 28, No. 4, pp. 477-491, July, 1990.
- [44] Sarabandi, K., "Scattering from variable resistive and impedance sheets", J. Electromag. Waves and Applics., vol. 4, no. 9, 1990.
- [45] Sarabandi, K., and T.B.A. Senior, "Low frequency scattering from cylindrical structures at oblique incidence", IEEE Trans. Geosci. Remote Sensing, vol. 28, no. 5, 879-885, Sept. 1990.
- [46] Sarabandi, K., "Simulation of a periodic dielectric corrugation with an equivalent anisotropic layer", J. Infrared and Millimeter Waves, vol. 11, no. 11, Nov. 1990.
- [47] Tavakoli, A., "Microwave Propagation Through a Man-Made Vegetation Canopy," Ph.D. dissertation, The University of Michigan, Ann Arbor, Michigan, 1991.
- [48] Ulaby, F. T., M. C. Dobson, and D. R. Brunfeldt, "Microwave Probe for In Situ Observations of Vegetation Dielectric", Proceedings of IEEE Instrumentation Measurement Technology Conference (IMTC/90), May 14-16, 1991, Atlanta, Georgia.
- [49] Kasischke, E. S., L. L. Bourgeau-Chavez, N. L. Christensen and M. C. Dobson, "The Relationship Between Aboveground Biomass and Radar Backscatter as Observed on Airborne SAR Imagery," Proceedings of the Third "Airborne Synthetic Aperture Radar (AIRSAR) Workshop, Jet Propulsion Laboratory, Pasadena, CA, May 20-24, 1991, JPL Publication 91-30, pp. 11 - 21, August 1, 1991.
- [50] DeRoo, R., Y. Kuga, M. C. Dobson, and F. T. Ulaby, "Bistatic Radar Scattering from Organic Debris of a Forest Floor," 1991 International Geoscience and Remote Sensing Symposium (IGARSS'91), IGARSS'91 Digest, IEEE91CH2971-0, Vol. 1, pp. 15-18, Espoo, Finland, June 3 -6, 1991.

- [51] Dobson, M. C., K. McDonald, and L. Pierce, "Diurnal Variation in Radar Backscatter from a Loblolly Pine Forest in Late Summer," 1991 International Geoscience and Remote Sensing Symposium (IGARSS'91), IGARSS'91 Digest, IEEE91CH2971-0, Vol. 3, pp. 1115, Espoo, Finland, June 3 - 6, 1991.
- [52] Sarabandi, K., Y. Oh, and F. Ulaby, "Polarimetric Radar Measurements of Bare Soil Surfaces at Microwave Frequencies", 1991 International Geoscience and Remote Sensing Symposium (IGARSS'91), IGARSS'91 Digest, IEEE91CH2971-0, Vol. 2, pp. 387-390, Espoo, Finland, June 3 - 6, 1991.
- [53] Ulaby, F.T., J. Stiles, K. Sarabandi and Y. Oh, "Model and Measurements of the Phase Statistics of the Scattering Matrix of Distributed Surfaces", 1991 International Geoscience and Remote Sensing Symposium (IGARSS'91), IGARSS'91 Digest, IEEE91CH2971-0, Vol. 2, pp. 401, Espoo, Finland, June 3 - 6, 1991.
- [54] Whitt, M.W. and F.T. Ulaby, "Observation and Modelling of Look-direction Effects in Microwave Backscatter from Corn", 1991 International Geoscience and Remote Sensing Symposium (IGARSS'91), IGARSS'91 Digest, IEEE91CH2971-0, Vol. 4, pp. 1887, Espoo, Finland, June 3 - 6, 1991.
- [55] DeRoo, R., Y. Kuga, F.T. Ulaby and M.C. Dobson, "Bistatic Radar Scattering from Organic Debris of a Forest Floor", 1991 International Geoscience and Remote Sensing Symposium (IGARSS'91), IGARSS'91 Digest, IEEE91CH2971-0, Vol. 1, pp. 15-18, Espoo, Finland, June 3 - 6, 1991.
- [56] Whitt, M.W. and F.T. Ulaby, "Modelling the Polarimetric Backscatter from Periodic Bare Soil Surfaces", 1991 International Geoscience and Remote Sensing Symposium (IGARSS'91), IGARSS'91 Digest, IEEE91CH2971-0, Vol. 2, pp. 391, Espoo, Finland, June 3 - 6, 1991.
- [57] Tavakoli, A., K. Sarabandi and F.T. Ulaby, "Horizontal Propagation through Periodic Vegetation Canopies", 1991 International Geoscience and Remote Sensing Symposium (IGARSS'91), IGARSS'91 Digest, IEEE91CH2971-0, Vol. 4, pp. 1879-1882, Espoo, Finland, June 3 - 6, 1991.
- [58] Dobson, M. C., R. DeLaSierra, and N. Christensen, "Spatial and Temporal Variation of the Microwave Dielectric Properties of Loblolly Pine Trunks," 1991 International Geoscience and Remote Sensing Symposium (IGARSS'91), IGARSS'91 Digest, IEEE91CH2971-0, Vol. 3, pp. 1107-1110, Espoo, Finland, June 3 - 6, 1991.
- [59] McDonald, K. C., M. C. Dobson and F. T. Ulaby, "Modeling Multi-Frequency Diurnal Backscatter from a Walnut Orchard," 1991 International Geoscience and Remote Sensing Symposium (IGARSS'91), IGARSS'91 Digest, IEEE91CH2971-0, Vol. 3, pp. 1125-1130, Espoo, Finland, June 3 - 6, 1991.
- [60] McDonald, K. and F.T. Ulaby, "Modeling Microwave Backscatter from Discontinuous Tree Canopies", Technical Report 026511-2-T, June 1991.
- [61] Whitt, M.W. and F.T. Ulaby, "An Efficient Method of Computing the Phase Matrix and the Extinction Matrix for Arbitrary Rayleigh Particles, 1991 North American Radio Science Meeting and International IEEE/AP-S Symposium, June 24-28, 1991, The University of Western Ontario, London, Ontario.

- [62] McDonald, K.C. and F.T. Ulaby, "Sensitivity of Tree Canopy Backscatter to variations in Open Canopy Parameters", 1991 North American Radio Science Meeting and International IEEE/AP-S Symposium, June 24-28, 1991, The University of Western Ontario, London, Ontario.
- [63] Tavakoli, A., K. Sarabandi, and F.T. Ulaby, "Microwave Propagation through Cultural Vegetation Canopies", Technical Report 026511-3-T, July 1991.
- [64] Whitt, M. and F.T. Ulaby, "Microwave Scattering from Periodic Row-Structured Vegetation", Technical Report 026511-4-T, June 1991.
- [65] Sarabandi, K., "Scattering from dielectric structures above impedance surfaces and resistive sheets," Proc. of Progress in Electromag. Research Symp., Boston, July 1991.
- [66] Sarabandi, K., A. Tavakoli and F.T. Ulaby, "Propagation in a Two Dimensional Periodic Random Media with Inhomogeneous Particle Distribution", Progress in Electromagnetics Research Symposium, 1-5 July, 1991, Cambridge, Mass.
- [67] Martens, S. N., S. L. Ustin, and J. M. Norman, "Measurement and Characterization of Tree Canopy Architecture, International Journal of Remote Sensing, Vol. 12, No. 7, pp. 1525-1546, July 1991.
- [68] Tavakoli, A., K. Sarabandi, F.T. Ulaby, "Horizontal Propagation Through Periodic Vegetation Canopies", IEEE Trans. Antennas & Propagation, Vol. 39, No. 7, pp. 1014-1023, July 1991.
- [69] Tavakoli, A., K. Sarabandi, and F. Ulaby, "Microwave Propagation Constant for a Vegetation Canopy at X-band", Submitted to Radio Science, August, 1991.
- [70] Ulaby, F.T., K. Sarabandi, and A. Nashashibi "Statistical Properties of the Mueller Matrix of Distributed Targets", Special Issue of IEE Proceedings Part-F, Vol. 139, No. 2, pp 136-146, April, 1992.
- [71] McDonald, K. C., "Modeling Microwave Backscatter From Tree Canopies," Ph.D. dissertation, The University of Michigan, Ann Arbor, Michigan, 1991.
- [72] Sarabandi, K., A. Tavakoli, and F.T. Ulaby, "Propagation in a Two Dimensional Periodic Random Medium with Inhomogeneous Particle Distribution", IEEE Trans. on Antennas and Propagation, Vol. 40, No. 10, pp. 1175-1186, October, 1992.
- [73] Way, JoBea, Eric Rignot, M. Craig Dobson and Kyle McDonald, "Monitoring the Environmental and Phenologic State of Alaskan Forests Using Synthetic Aperture Radar," submitted to the 1991 Annual Meeting of the Ecological Society of America, August 4-8, 1991, San Antonio, Texas.
- [74] Haney, Eric, Norman Christensen, Eric Kasischke, and M. Craig Dobson, "The Impact of Variations in Tree Architecture and Density on Synthetic Aperture Radar Signatures of Loblolly Pine Forests in the Duke University Research Forest," 1991 Annual Meeting of the Ecological Society of America, August 4-8, 1991, San Antonio, Texas.

- [75] Whitt, M. W., "Microwave Scattering from Periodic Row-Structured Vegetation," Ph.D. dissertation, The University of Michigan, Ann Arbor, Michigan, 1991.
- [76] Way, J. B., M. Schier, M. C. Dobson, D. Gates, F. T. Ulaby, S. L. Ustin, V. Vanderbuilt, J. Weber, J. Paris, E. Kasischke, R. Lang and J. Norman, "Diurnal Change in Trees as Observed by Optical and Microwave Sensors: The EOS Synergism Study," IEEE Transactions Geoscience and Remote Sensing, Vol. 29, No. 6, pp. 807-821, November, 1991.
- [77] Casey, D. and J. B. Way, "Orbit Selection for the EOS Mission and Its Synergism Implications," IEEE Trans. Geoscience and Remote Sensing, Vol.29, No. 6, pp. 822-835, November 1991.
- [78] Gates, D. M., "Water Relations of Forest Trees," IEEE Trans. Geoscience and Remote Sensing, Vol.29, No. 6, pp. 836-842, November 1991.
- [79] Ustin, S. L., S. N. Martens, and V. C. Vanderbuilt, "Canopy Architecture of a Walnut Orchard," IEEE Trans. Geoscience and Remote Sensing, Vol.29, No. 6, pp. 843-851, November 1991.
- [80] McDonald, K. C., M. C. Dobson, and F. T. Ulaby, "Modeling Multi-Frequency Diurnal Backscatter from a Walnut Orchard," IEEE Trans. Geoscience and Remote Sensing, Vol.29, No. 6, pp. 852-863, November 1991.
- [81] Weber, J. A. and S. L. Ustin, "Diurnal Water Relations of Walnut Trees: Implications for Remote Sensing," IEEE Trans. Geoscience and Remote Sensing, Vol.29, No. 6, pp. 864-874, November 1991.
- [82] Vanderbuilt, V. C., S. L. Ustin, K. M. Berger, W. F. Caldwell, J. A. Clark, B. D. Ganapol, E. S. Kasischke, S. N. Martens, R. E. Pettigrew, and R. A. Rousseau, "Biophysical Information in Asymmetric and Symmetric Diurnal Bidirectional Canopy Reflectance," IEEE Trans. Geoscience and Remote Sensing, Vol.29, No. 6, pp. 875-889, November 1991.
- [83] McDonald, K.C. and F.T. Ulaby, "Radiative Transfer Modeling of Discontinuous Tree Canopies at Microwave Frequencies", submitted for publication in International Journal of Remote Sensing, December, 1991.
- [84] Whitt, M.W. and F.T. Ulaby, "Coherent Stalk Effects in Microwave Backscatter from Corn", submitted for publication, January, 1992.
- [85] Sarabandi, K., "Scattering from dielectric structures above impedance surfaces and resistive sheets," IEEE Trans. Antennas Propagat., vol. 40, no. 1, Jan. 1992.
- [86] Oh, Y., K. Sarabandi, and F.T. Ulaby, "An Empirical Model and an Inversion Technique for Radar Scattering from Bare Soil Surfaces", IEEE Transactions on Geosci. and Remote Sensing, Vol. 30, pp. 370-381, March, 1992.
- [87] Dobson, M. C., F. T. Ulaby, T. LeToan, A. Beaudoin, E. S. Kasischke, and N. Christensen, "Dependence of Radar Backscatter on Conifer Forest Biomass," IEEE Trans. Geoscience and Remote Sensing, Vol. 30, No. 2, pp. 412-415, March, 1992.

- [88] Whitt, M.W. and F.T. Ulaby, "A Microwave Scattering Model for Periodic Row-Structured Vegetation", submitted for publication, May, 1992.
- [89] Sarabandi, K., "Derivation of phase statistics of distributed targets from the Mueller matrix," Radio Sci., vol. 27, no. 5, pp 553-560, 1992.
- [90] Sarabandi, K., Y. Oh, and F.T. Ulaby, "An Empirical Model for Phase Difference Statistics of Rough Surfaces", URSI Specialist Meeting on Microwave Signatures, Innsbruck, Austria, July 1-3, 1992.
- [91] Sarabandi, K., Y. Oh, and F.T. Ulaby, "Scatterometer Measurement of Differential Mueller Matrix of Distributed Targets," 1992 International Geoscience and Remote Sensing Symposium (IGARSS'92), Houston, Texas, May 26 - 29, 1992.
- [92] Sarabandi, K., P.F. Polatin, F.T. Ulaby, "Monte Carlo Simulation of Scattering from a Layer of Vertical Cylinders", 1992 International Geoscience and Remote Sensing Symposium (IGARSS'92), Houston, Texas, May 26 - 29, 1992.
- [93] Dobson, M. C. and E. Wilcox, "Role of Crown Structure in Radar Backscatter Dependence on Forest Biomass," 1992 International Geoscience and Remote Sensing Symposium (IGARSS'92), Houston, Texas, May 26 - 29, 1992.
- [94] Polatin, P.F., K. Sarabandi, and F.T. Ulaby, "An Iterative Method for Inversion of the Polarimetric Radar Response of Vegetation Canopies ", 1992 International Geoscience and Remote Sensing Symposium (IGARSS'92), Houston, Texas, May 26 - 29, 1992.
- [95] Pierce, L.E., K. Sarabandi, J.M. Stiles, and F.T. Ulaby, "Application of an Artificial Neural Network in a Canopy Scattering Model Inversion", 1992 International Geoscience and Remote Sensing Symposium (IGARSS'92), Houston, Texas, May 26 - 29, 1992.
- [96] Nashashibi, A., K. Sarabandi, and F.T. Ulaby, "The Effect of Azimuthal Asymmetry of Particle Orientation on the Phase Statistics of Distributed Targets", 1992 International Geoscience and Remote Sensing Symposium (IGARSS'92), Houston, Texas, May 26 - 29, 1992.
- [97] Stiles, J.M., K. Sarabandi and F.T. Ulaby, "Polarimetric Scattering Model for Grassy Terrain", APS Meeting, Adelaide , Chicago, 1992.
- [98] Polatin, P.F., K. Sarabandi and F.T. Ulaby, "An Iterative Inversion Algorithm with Application to the Polarimetric Radar Response of Vegetation Canopies", submitted for publication, July 1992.
- [99] Ulaby, F.T., Y. Oh, and K. Sarabandi, "Inversion Model for Retrieving Soil Surface Moisture and Roughness from Polarimetric Radar Observations", URSI Specialist Meeting on Microwave Signatures, Innsbruck, Austria, July 1-3, 1992.
- [100] Sarabandi, K., "Derivation of phase statistics from Mueller matrix," URSI Specialist Meeting on Microwave Signatures, Innsbruck, Austria, July 1-3, 1992.

- [101] Pierce, L.E., K. Sarabandi and F.T. Ulaby, "Application of an Artificial Neural Network in Canopy Scattering Inversion", submitted to IEEE Transactions on Geoscience and Remote Sensing, August, 1992.
- [102] Way, J. B., R. Oren, R. Kwok, E. Rignot, G. Bonan, K. McDonald, C. Dobson, L. Viereck and J. E. Roth, "Monitoring Temporal Change and Species in Alaskan Forests Using AIRSAR and ERS-1 Data," accepted by IEEE Transactions Geoscience and Remote Sensing, October, 1992.
- [103] Sarabandi, K., Y. Oh, and F.T. Ulaby, "Measurement and Calibration of Differential Mueller Matrix of Distributed Targets", IEEE Transactions on Antennas and Propagation, Vol. 40, No. 12, pp. 1524-1532, December, 1992.
- [104] Ulaby, F. T. and M. C. Dobson, "Radar Response of Vegetation: An Overview," Third Spaceborne Imaging Radar Symposium, Jan. 18 - 21, 1993, Jet Propulsion Laboratory, Pasadena, CA.
- [105] Dobson, M. C., K. McDonald, F. T. Ulaby, E. S. Kasischke, and J. B. Way, "Modeling the Effects of Temperature on Microwave Extinction and Backscatter from a Boreal Forest in Winter", submitted to IEEE Transactions Geoscience and Remote Sensing, January, 1993.
- [106] Sarabandi, K., P.F. Polatin and F.T. Ulaby, "Monte Carlo Simulation of Scattering from a Layer of Vertical Cylinders", accepted for publication in IEEE Trans. on Antennas and Propagation, February, 1993.
- [107] Stiles, J.M., K. Sarabandi and F.T. Ulaby, "Microwave Scattering Model for Grass Blade Structures", accepted for publication in IEEE Trans. Geoscience and Remote Sensing, April, 1993.
- [108] Sarabandi, K., Y. Oh, and F.T. Ulaby, "A Numerical Simulation of Scattering From Inhomogeneous Dielectric Random Surfaces", submitted for publication in AP, April, 1993
- [109] DeRoo, R., and F.T. Ulaby, "Bistatic Spectral Scattering from Rough Dielectric Surfaces", IEEE Trans. Antennas Propagat., submitted for publication , April, 1993.
- [110] Sarabandi, K., and P.F. Polatin, "Electromagnetic scattering from two adjacent objects," IEEE Trans. Antennas Propagat., submitted for publication May, 1993.
- [111] Sarabandi, K., Y. Oh and F.T. Ulaby, "A Numerical Simulation of Scattering from an Inhomogeneous Random Surface", International Antennas and Propagation Symposium, June, 1993, Ann Arbor, Michigan.
- [112] Oh, Y., K. Sarabandi and F. Ulaby, "Re-Examination of the Kirchhoff Approximation for Scattering from a Rough Surface", International Antennas and Propagation Symposium, June, 1993, Ann Arbor, Michigan.
- [113] Stiles, J.M., K. Sarabandi, T.B.A. Senior, F.T. Ulaby, "Scattering formulation for long thin cylinders of arbitrary cross-section and Electrical length", Progress in Electromagnetics Research Symposium, July, 1993, Pasadena, California.

- [114] Sarabandi, K., P.F. Polatin, ``Electromagnetic scattering from two adjacent objects," Progress in Electromagnetics Research Symposium, July, 1993, Pasadena, California.
- [115] Pierce, L. E., K. Sarabandi, F. T. Ulaby and M. C. Dobson, "Knowledge-Based Classification of SAR Images," 1993 International Geoscience and Remote Sensing Symposium (IGARSS'93), August 18 - 21, 1993, Kogakuin University, Tokyo, Japan.
- [116] Pierce, L. E., M. C. Dobson, E. Wilcox and F. T. Ulaby, "Artificial Neural Network Inversion of Tree Canopy Parameters in the Presence of Diversity," 1993 International Geoscience and Remote Sensing Symposium (IGARSS'93), August 18 - 21, 1993, Kogakuin University, Tokyo, Japan.
- [117] DeRoo, R.D. and F.T. Ulaby, "Experimental Observations of the Forward Scattering of Microwaves from a Rough Dielectric Surface", 1993 International Geoscience and Remote Sensing Symposium (IGARSS'93), August 18 - 21, 1993, Kogakuin University, Tokyo, Japan.
- [118] Oh, Y., K. Sarabandi and F.T. Ulaby, "An Empirical Model for Phase Differences Statistics of Rough Surfaces", 1993 International Geoscience and Remote Sensing Symposium (IGARSS'93), August 18 - 21, 1993, Kogakuin University, Tokyo, Japan.
- [119] Dobson, M. C., E. Wilcox and F. T. Ulaby, "Effects of Forest Structure on Radar Response to Forest Biomass," 1993 International Geoscience and Remote Sensing Symposium (IGARSS'93), August 18 - 21, 1993, Kogakuin University, Tokyo, Japan.